Ref: IMOCSCG019
South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-professionals/medicines/sel-imoc/sel-imoc-shared-care-professionals/medicines/sel-imoc-shared-care-profess



SHARED CARE PRESCRIBING GUIDELINE for the prescribing and monitoring of stimulant therapy and anti-cataplectic agents in the management of Narcolepsy (+/- Cataplexy) and Idiopathic Hypersomnia in adults

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

SHARED CARE PROCESS FLOWCHART

Specialist clinician completes Shared Care Request Letter (Appendix 1) and sends to patient's GP via email.



GP considers shared care request, taking into account the following:

- Is the patient's condition predictable or stable?
- Whether they have the relevant knowledge, skills and access to equipment to allow them to monitor treatment as indicated in this shared care prescribing guideline?
- Whether they have been provided with relevant clinical details including monitoring data?





If YES to all the above, and after reading this shared care guideline then it is appropriate for GP to accept prescribing responsibility



If NO to any of these questions, GP should contact the requesting consultant or the local primary care Medicines Optimisation Team within 2 weeks of receipt to discuss



Complete Shared Care Agreement Letter (Appendix 2) and email back to the requesting clinician within 2 weeks of receipt

Complete Shared Care Refusal Letter (Appendix 3) and email back to the requesting clinician

NOTES

There may be implications for the patient where invitation to share care is declined. For example, the patient may need to be changed to an alternative treatment regimen. It would not normally be expected that shared care prescribing would be declined on the basis of cost.

Sharing of care assumes communication between the specialist, GP and patient. The intention to share care should be explained to the patient by the doctor initiating treatment. It is important that patients are consulted about treatment and are in agreement with it.

Prescribing should follow requirements in the <u>South East London Interface Prescribing Policy</u>. The doctor who prescribes the medication legally assumes clinical responsibility for the drug and the consequences of its use. The patient's best interests are always paramount.

If the GP is not confident to undertake these roles, then he or she is under no obligation to do so. In such an event, the total clinical responsibility for the patient for the diagnosed condition remains with the specialist. If a specialist asks the GP to prescribe this drug, the GP should reply to this request as soon as practicable (within 2 weeks).

Idiopathic Hypersomnia in adults - stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-agreements/

1. AREAS OF RESPONSIBILITY

It is the responsibility of the specialist team to work with the Primary Care Lead to support GPs with drug monitoring, including consideration of patient recall systems where appropriate, and to advise on long-term stock issues where these become apparent.

Consultant / Specialist team responsibilities

- Ensure the stimulant therapy or anti-cataplectic agent is appropriately initiated e.g. no contraindications, cautions,
- Ensure prescribing fits local or national agreement for use of the drug; off-label use of the drug should be discussed with the patient
- Counsel the patient on the risks and benefits of treatment, discuss potential adverse effects and any practical issues related to the use of stimulant therapy or anti-cataplectic agents
- Provide the patient with written information on the drug where available
- Provide the patient with information on self-management and advise them to contact secondary care specialist if they experience a deterioration of their disease management
- Undertake baseline investigations and initial monitoring according to local protocol
- Advise the patient to sign up to access their GP record online or via a mobile phone app where available; if this is not possible or suitable for the patient then ensure they are provided with a medicines monitoring record and advised of the importance of ongoing monitoring
- Advise the patient to be up to date with their routine vaccinations
- Signpost the patient to additional support services, such as charities and patient groups, where available and appropriate
- Report any suspected adverse effects to the MHRA: https://yellowcard.mhra.gov.uk/
- Discuss shared care with patient and obtain patient agreement to request shared care
- Prescribe treatment for at least the first 3 months or until the patient is considered stable and shared care is agreed with GP; ongoing responsibility for prescribing stimulant therapy or anti-cataplectic agent will remain with the specialist team until this has been taken over by the GP
- If the GP does not agree to shared care continue to prescribe and monitor the stimulant therapy or anti-cataplectic agents
- When shared care is agreed:
 - Review patient at the request of GP should any problems arise (monitoring/ side-effects / lack of efficacy/ noncompliance)
 - Arrange routine follow up (usually at least once per year)
 - Communicate (within 2 weeks) with the GP if treatment is changed or stopped; if urgent communicate the change within 48 hours
 - Confirm the patients monitoring schedule at each appointment.

General Practitioner responsibilities

- To respond to shared care proposal within 2 weeks of receipt
- Provide ongoing prescriptions and adjust dose as advised by the specialist; annotate the prescription 'as per shared care guideline'
- Inform the patient of the monitoring arrangement required
- Encourage patient to register to access their record online and ensure this is updated; if the patient does not have online access enter blood results in patient held medicines monitoring record (for replacement or renewal of patient held medicines monitoring records contact the secondary care team)
- Undertake monitoring as outlined in the monitoring table
- To comment on the results of any monitoring undertaken in primary care to make the results and any impact of these clear to the patient
- Refer to the 'actions to be taken in primary care' page in the shared care guideline and follow the action specified
- Refer back to secondary care if the patient's condition deteriorates
- Stop treatment on the advice of secondary care or immediately if an urgent need to stop treatment arises
- Report any suspected adverse effects to the MHRA via the Yellow Card scheme: https://yellowcard.mhra.gov.uk/
- Refer the patient back to secondary care if they are planning a family (males and females).

Patient's / Carer's responsibilities

- Read pre-treatment information leaflets when provided by secondary care
- Attend scheduled appointments in primary and secondary care
- Contact the secondary or primary care team if unclear on any aspect of the treatment
- Report concerns about side effects to a healthcare professional
- Report any plans to start a family/ breastfeed to primary or secondary care
- Report any new or worsening symptoms to primary or secondary care
- Inform primary or secondary care or community pharmacist of any other medication being taken, including over the counter products or herbal remedies
- Take medicines as agreed and try to ensure no doses are missed
- Keep contact details up to date with both primary and secondary care

Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

Idiopathic Hypersomnia in adults - stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-agreements/

To arrange and attend all regular monitoring required and to confirm with the specialist or GP team that the results of these are satisfactory to continue the medication (contact can be via telephone to the admin team or patient having online access to their results in cases where confirmation is provided alongside the results that they are satisfactory)

2. CLINICAL INFORMATION

NOTE: The information here is not exhaustive. Please also consult the current Summary of Product Characteristics (SPC) for the stimulant therapy and anti-cataplectic agents listed below prior to prescribing for up to date prescribing information, including detailed information on adverse effects, drug interactions, cautions and contraindications (available via www.medicines.org.uk)

Background

Narcolepsy (+/- cataplexy) and idiopathic hypersomnia are both long-term and debilitating sleep conditions which are similar in their clinical presentation, differ in their diagnosis but may be managed in a parallel manner.

Narcolepsy is a long-term condition that causes excessive sleepiness during the day and may also disrupt your sleep at night. You can also have sleep attacks where you fall asleep at inappropriate times during the day without any warning. The Epworth Sleepiness Scale is a questionnaire intended to measure daytime sleepiness. Narcolepsy is generally associated with an ESS of >12, even after adequate night-time sleep.

Cataplexy is a condition associated with narcolepsy that results in sudden muscle weakness triggered by strong emotions such as laughter, anger, fright or surprise. Muscle weakness can vary in severity and cataplexy attacks can differ in both nature and duration.

Idiopathic Hypersomnia (IH) is a sleep disorder in which a person is excessively sleepy during the day and has great difficulty being awakened from sleep. Idiopathic means there is no clear cause. IH is similar to narcolepsy in that you are extremely sleepy but also different from narcolepsy because IH does not usually involve suddenly falling asleep (sleep attacks) or losing muscle control due to strong emotions (cataplexy). Furthermore, unlike narcolepsy, naps in idiopathic hypersomnia are usually not refreshing.

The purpose of this document is to demonstrate the clinical use of stimulant and anti-cataplectic agents in the treatment of narcolepsy (+/- cataplexy) and idiopathic hypersomnia in adult patients. It is not within the scope of this document to provide guidance on diagnosis of this condition.

Indications

Note if indication is unlicensed or not

The table below indicates the stimulant therapy and anti-cataplectic agents used in the treatment of narcolepsy (+/- cataplexy) and idiopathic hypersomnia in adult patients O = 'off-label' but considered routine treatment option

X = unlicensed and not currently considered a routine option			
	Narcolepsy	Narcolepsy with Cataplexy	Idiopathic Hypersomnia
Stimulant agents			
Modafinil	Licensed	Licensed	0
Methylphenidate XL	0	0	0
Methylphenidate IR	0	0	0
Dexamfetamine	Licensed	Licensed	0
Anti-cataplectic agents			
Sodium oxybate	0	Licensed	X
Venlafaxine XL/IR	X	0	X
Clomipramine	Χ	Licensed	X
Fluoxetine	X	0	X
The agent below is red listed and is NOT included in this shared care arrangement. Prescribing will remain with the sleep centre.			
Pitolisant (NOT FOR SHARED CARE)	Licensed	Licensed	0
Solriamfetol (NOT FOR SHARED CARE)	Licensed	Licensed	X

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Place in Therapy Indicate what drugs should have been tried before this drug is considered Locally agreed off-label use Including supporting information	See the following clinical pathways for further information: • Pathway for the pharmacological management of excessive daytime sleepiness due to narcolepsy • Pathway for the pharmacological management of cataplexy associated with narcolepsy See the South East London Joint Medicines Formulary for more information	
Initiation and ongoing dose regime Note: • Transfer of monitoring and prescribing to primary care is normally after the patient's dose has been optimized and with satisfactory investigation results for at least 12 weeks. • The duration of treatment & frequency of review will be determined by the specialist, based on clinical response and tolerability. • All dose or formulation adjustments will be the responsibility of the initiating specialist unless directions have been discussed and agreed with the primary care clinician. • Termination of treatment will be the responsibility of the specialist.	Initial stabilisation: (The loading period must be prescribed by the initiating specialist) This will be for at least 3 months Some patients may take longer for their dose to stabilise, and in these cases, prescribing will remain with the initiating specialist until stabilized Maintenance dose (following initial stabilisation): After the initial stabiliation, GPs can be asked to continue prescribing treatment as per SE London shared care agreement (see appendix 1 below) Conditions requiring dose adjustment Sub-therapeutic Mild side-effects i.e. headache Some patients may have more individualised parameters set out by their secondary care specialist which fall outside the normal range; these should be communicated to primary care in writing. Duration of treatment Ongoing provided there is benefit, no reported intolerance and/or adverse	
Pharmaceutical aspects	effects. Route of administration Formulation Administration See appendix 4 below Administration details Other important See appendix 4 below See appendix 4 below	
Baseline investigations, initial monitoring and ongoing monitoring to be undertaken by specialist	Baseline investigations: Prior to initiation the following pre-treatment investigations will be undertaken in secondary care: Height & weight Blood pressure Heart rate Epworth Sleep Score Pregnancy test (if required) ECG (for initiation of modafinil, methylphenidate and dexamfetamine). Cataplexy severity (for initiation of venlafaxine, clomipramine, fluoxetine and sodium oxybate) Oximetry and mood (for initiation of sodium oxybate) In addition, the initiating secondary care specialist is responsible for: Documenting pre-treatment disease status as appropriate by indication Considering contraindications to treatment, co-morbidities or patient factors that would influence drug choice Counselling the patient on the risks and benefits of treatment including potential side effects, the need for ongoing monitoring and disease management To initiate, stabilise and supply treatment for at least 3 months.	

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Ongoing monitoring requirements to be undertaken by primary care Adverse effects and	anti-cataplectic agents, suc dose. Initial monitoring Monitoring at baseline and du only once the patient is optimi	cal issues related to the use of stimulant +/- h as administration, storage and maximum ring initiation is the responsibility of the specialist, ised on the chosen medication with no anticipated nmediate future will prescribing and monitoring be Frequency See appendix 4 below Action for GP
management Any serious adverse reactions should be reported to the MHRA via the Yellow Care scheme www.mhra.gov.uk/yellowcard	See appendix 4 below	Inform the named consultant of any reported adverse effects.
Advice to patients and carers The specialist will counsel the patient with regard to the benefits and risks of treatment and will provide the patient with any relevant information and advice, including patient information leaflets on individual medicines. Criteria for stopping treatment e.g. poor response, adverse effects requiring cessation	The patient should be advised to report any of the following signs or symptoms to their GP without delay: Rash develops. Onset of chest pain. Palpitations, breathlessness of unknown cause. Changes in mood, behaviour or thinking, suicidal thoughts. Stopping criteria (for all stimulant therapy and anti-cataplectic agents) Inform named consultant urgently. The sleep centre will advise appropriate weaning regimen. Failure to respond to treatment or adverse effects necessitating withdrawal. Pregnancy Patient request Rash develops, onset of chest pain, palpitations, breathlessness of unknown cause, and changes in mood, behaviour or thinking, suicidal	
Follow up arrangements e.g. frequency of specialist clinic attendance	thoughts. Request patient seen earlier if condition deterioration or adverse effects experienced between appointments It is the primary care prescriber's responsibility to ensure patients adhere to the monitoring schedule. It should be clearly communicated to the patient how often they are required to attend. Concerns that the patient is unable to adhere to the monitoring schedule should be discussed with the secondary care team. Patients must be informed that they will be unable to continue the medication unless they adhere to the monitoring requirements.	
Pregnancy, paternal exposure, and breast feeding It is the responsibility of the specialist to provide advice on the need for contraception to male and female patients on initiation and at each review but the ongoing responsibility for providing this advice rests with both the GP and the specialist.	Pregnancy: See appendix 4 below Breastfeeding: See appendix 4 below For Specialist	

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

	Please complete the modafinil consent form (internal GSTT sleep centre consent form)		
Additional information	Where patient care is transferred from one specialist service or GP practice to another, a new shared care agreement must be completed.		
	Some patients may have more individualised parameters set out by their secondary care specialist which fall outside the normal range; these should be communicated to primary care in writing.		
Evidence base for treatment	Summary of Product Characteristics. Available online at:		
and key references	https://www.medicines.org.uk/emc#gref		
Include hyperlinks to original sources and	2. British National Formulary. Available online at: https://bnf.nice.org.uk/		
access dates	3. Department of Health. Green Book. Available online:		
	https://www.gov.uk/government/collections/immunisation-against-		
	infectious-disease-the-green-book#the-green-book		
	 Department of Health. MHRA alert. Available online: https://www.gov.uk/drug-safety-update/modafinil-provigil-increased-risk-of-congenital-malformations-if-used-during-pregnancy 		
To be read in conjunction with	 South East London Joint Medicines Formulary Formulary 		
the following documents	 Pathway for the pharmacological management of excessive daytime 		
	sleepiness due to narcolepsy		
	 Pathway for the pharmacological management of cataplexy associated 		
	with narcolepsy		
Local arrangements for referral	As per shared care process flow chart.		
Define the referral procedure from hospital to primary care prescriber & route of return should the patient's condition change.	Clinic letter/email request to GP for shared care consideration. Practice letter/email from GP to secondary care.		

3. COMMUNICATION AND SUPPORT

Guy's and St. Thomas' Hospital switchboard: 0207 188 7188		
Consultant/specialist team Rex Muza (Clinical Lead)	Tel: 0207 188 3430 Email: gst-tr.gsttsleepreferrals@nhs.net	
Medication – Prescribing advice, interactions, availability of medicines Jasvinder Kaler Singh	Tel: 0207 188 3430 Email: gst-tr.thesleeppharmacistgstt@nhs.net	
Medicines Information Guy's Hospital Medicines Information Department If you have any questions or concerns about these medicines, please call our helpline.	Tel: 0207 188 3855/3853 Email: gstt.medicinesinformation@nhs.net	
Guy's Hospital Sleep Disorder Centre Open Mon-Fri 9am-5pm	Tel: 0207188 3430 Email: gst-tr.gsttsleepreferrals@nhs.net	
Language and accessible support services If you need an interpreter or information about the pharmacological management of these conditions in a different language or format, please get in touch.	Tel: 020 7188 8815 Email: gstt.languagesupport@nhs.net	
NHS 111 Offers medical help and advice from fully trained advisers supported by experienced nurses and paramedics. Available over the phone 24 hours a day.	Tel: 111	
NHS Choices Provides online information and guidance on all aspects of health and healthcare.	Website: www.nhs.uk	

Idiopathic Hypersomnia in adults - stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Appendix 1: Shared Care Request letter (Specialist to Primary Care Prescriber)

Dear [insert Primary Care Prescriber's name]

Patient name: [insert patient's name]
Date of birth: [insert date of birth]
NHS Number: [insert NHS Number]
Diagnosis: [insert diagnosis]

As per the agreed South East London shared care prescribing guideline for [insert medicine name] for the treatment of [insert indication], this patient is now suitable for prescribing to move to primary care.

The patient fulfils criteria for shared care and I am therefore requesting your agreement to participate in shared care. Where baseline investigations are set out in the shared care protocol, I have carried these out.

I can confirm that the following has happened with regard to this treatment:

[Shared care can only be considered if the following requirements have been met. Please complete all parts of the right hand column to confirm this]	Specialist to complete
The patient has been initiated on this therapy and has been on an optimised dose for the following period of time:	weeks/months
Baseline investigation and monitoring as set out in the shared care documents have been completed and were satisfactory	Yes □
The condition being treated has a predictable course of progression and the patient can be suitably maintained by primary care	Yes □
The risks and benefits of treatment have been explained to the patient	Yes □
A contraceptive check for this patient has been completed within the last months/week	Yes, Dated: □ N/A □
The roles of the specialist/specialist team/ Primary Care Prescriber / Patient and pharmacist have been explained and agreed	Yes □
The patient has agreed to this shared care arrangement, understands the need for ongoing monitoring, and has agreed to attend all necessary appointments	Yes □
I have enclosed a copy of the shared care protocol which covers this treatment/the SCP can be found here (insert electronic/ web link)	Yes □
I have included with the letter copies of the information the patient has received	Yes □
I have provided the patient with sufficient medication to last until	
I have arranged a follow up with this patient in the following timeframe e.g. within 3 months / 6 months (please specify)	

Treatment was started on [insert date started] and the current dose is [insert dose and frequency].

If you are in agreement, please undertake monitoring and treatment from [insert date] NB: date must be at least 1 month from initiation of treatment.

The next blood monitoring is due on [insert date] and should be continued in line with the shared care guideline.

Please could you reply to this request for shared care and initiation of the suggested medication to either accept or decline within 14 days.

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South 8
East London Integrated Care System: SEL Integrated Care Board (covering the boroughs of Bexley/Bromley/Greenwich/
Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

Identifier

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults - stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Appendix 2: Shared Care Agreement Letter (Primary Care Prescriber to Specialist)

Dear [insert Doctor's name]			
Patient	[insert Patient's name]		
NHS Number	[insert NHS Number]		

[insert patient's date of birth and/oraddress]

Primary Care Prescriber Response

Thank you for your request for me to accept prescribing responsibility for this patient under a shared care agreement and

mulant therapy			
Modafinil	Methylphenid	ate	Dexamfetamine
ti-cataplectic agents			
Venlafaxine XL	Venlafaxine	IR /	Clomipramine
Fluoxetine	Sodium oxyb	ate	
	\square		
	the primary care team agree to s; annotate the prescription 'as pe		
care guideline Report any concerns e.g. adv Refer to abnormal blood test the action specified Refer the patient back to secure Refer the patient back to secure unexpected pregnancy Encourage the patient to sign available or if the patient doe blood test results are docume Stop treatment on the advice	y secondary care and undertake a verse effects, inefficacy and non-commonitoring or adverse effect table ondary care if the patient's mediciondary care if they are planning an up to access their GP online sers not wish to have online access ented in the patient's medicine more of the secondary care or immediated, please document reason(s	compliance to sect in the shared car al condition deter a family (males ar vices/ via mobile then ensure that conitoring record ately if an urgent	condary care for advice are guideline and follow iorates ad females) or report an app if available, if not any changes in treatment and
J	,	, <u>.</u>	
ompleted by: Job Title:	Signature:	Da	to.
ompleted by, Job Title.	Signature.	Da	ie.
n confirm that I am willing to	take on this responsibility from [insert datel and v	will complete the monitoring
he shared care protocol for th	nis medicine/condition.		
•	•		
mary Care Prescriber signature		_	ate:

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults - stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Primary Care Prescriber address/practice stamp:

Appendix 3: Shared Care Refusal Letter (Primary Care Prescriber to Specialist)

Re:

Patient [insert Patient's name]

NHS Number [insert NHS Number]

Identifier [insert patient's date of birth and/oraddress]

Thank you for your request for me to accept prescribing responsibility for this patient.

In the interest of patient safety, the local NHS in South East London have classified [insert medicine name] as a Shared Care medicine, and requires a number of conditions to be met before transfer can be made to primary care.

I regret to inform you that in this instance I am unable to take on responsibility due to the following:

		Tick which
		apply
1.	The prescriber does not feel clinically confident in managing this individual patient's condition, and there is a sound clinical basis for refusing to accept shared care	
	As the patients primary care prescriber I do not feel clinically confident to manage this patient's condition because [insert reason]. I have consulted with other primary care prescribers in my practice who support my decision. This is not an issue which would be resolved through adequate and appropriate training of prescribers within my practice.	
	I have discussed my decision with the patient and request that prescribing for this individual remain with you as the specialist, due to the sound clinical basis given above.	
2.	The medicine or condition does not fall within the criteria defining suitability for inclusion in a shared care arrangement	
	As the medicine requested to be prescribed is not included on the national list of shared care drugs as identified by RMOC (Regional Medicines Optimisation Committees) or is not a locally agreed shared care medicine I am unable to accept clinical responsibility for prescribing this medication at this time.	
	Until this medicine is identified either nationally or locally as requiring shared care the responsibility for providing this patient with their medication remains with you	
3.	A minimum duration of supply by the initiating clinician	
	As the patient has not had the minimum supply of medication to be provided by the initiating specialist I am unable to take clinical responsibility for prescribing this medication at this time. Therefore can you please contact the patient as soon as possible in order to provide them with the medication that you have recommended.	
	Until the patient has had the appropriate length of supply the responsibility for providing the patient with their medication remains with you.	
4.	Initiation and optimisation by the initiating specialist	
	As the patient has not been optimised on this medication I am unable to take clinical responsibility for prescribing this medication at this time. Therefore can you please contact the patient as soon as possible in order to provide them with the medication that you have recommended.	
	Until the patient is optimised on this medication the responsibility for providing the patient with their medication remains with you.	

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

5.	Shared Care Protocol not received	
	As legal responsibility for clinical care lies with the clinician who signs the prescription, I need to ensure that I am in possession of sufficient clinical information for me to be confident to prescribe this treatment for my patient and it is clear where each of our responsibilities lie to ensure the patient is safely managed.	
	For this reason I am unable to take clinical responsibility for prescribing this medication at this time, therefore would you please contact the patient as soon as possible in order to provide them with the medication that you have recommended.	
	Until I receive the appropriate SCP, responsibility for providing the patient with their medication remains with you.	
6.	Other (Primary Care Prescriber to complete if there are other reasons why shared care cannot be accepted. NB: Capacity issues to be discussed with local primary care Medicines Optimisation Team prior to returning this form)	

I would be willing to consider prescribing for this patient once the above criteria have been met for this treatment.

NHS England 'Responsibility for prescribing between Primary & Secondary/Tertiary care' guidance (2018) states that "when decisions are made to transfer clinical and prescribing responsibility for a patient between care settings, it is of the utmost importance that the GP feels clinically competent to prescribe the necessary medicines. It is therefore essential that a transfer involving medicines with which GPs would not normally be familiar should not take place without full local agreement, and the dissemination of sufficient, up-to-date information to individual GPs." In this case we would also see the term GP being interchangeable with the term Primary Care Prescriber.

Please do not hesitate to contact me if you wish to discuss any aspect of my letter in more detail and I hope to receive more information regarding this shared care agreement as soon as possible

Yours sincerely	
Primary Care Prescriber signature:	Date:
Primary Care Prescriber address/practice stamp:	

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc/sel-imoc-shared-care-agreements/

Appendix 4 - Drug monitoring summary

Modafinil – refer to pathway for place in therapy (see links on page 5)

	or narcolepsy, modafinil can help to control symptoms.				
Route, Dose, Duration	Monitoring Undertaken by Specialist before requesting shared care	Ongoing monitoring to be undertaken by GP	Stopping Criteria	Monitoring following dose changes	Follow Up
Oral: modafinil 50-100mg/day gradually increasing over 4 to 6 weeks to maintenance dose of 400mg/day. Renal impairment GFR<10 Start at 50% normal dose and increase according to response. Hepatic impairment Reduce dose by half in patients with severe hepatic impairment. Older people Limited data available but potential for lower clearance and increased systemic exposure, it is recommended that patients ≥ 65 years of age commence therapy at 100 mg daily. Duration of Treatment Indefinitely if patient is responding well to treatment and in absence of significant side effects.	Baseline Blood pressure, heart rate, pregnancy test if necessary, Epworth Sleep Score. Ongoing at 6 and 12 monthly clinic appointments Blood pressure, Epworth Sleep Score Ask patient about any rashes or change in mood and behaviour at each visit. Ask patient about side effects such as headache, chest pain. Hormonal contraceptives may be less effective when used with modafinil and therefore are not recommended. Modafinil consent form to be completed. Completion of modafinil consent form (internal GSTT sleep centre consent form) Please refer to this guidance for appropriate contraceptive methods: MHRA advice	Blood pressure, ECG and heart rate annually. Results of concern may be discussed with the Sleep Centre. (Please see page 8 for contact details) Adverse reactions and inform the named consultant if concerns that the patient may be misusing the medication.	Failure to respond to treatment or adverse effects necessitating withdrawal. Pregnancy Patient request STOP modafinil if rash develops, onset of chest pain, palpitations, breathlessness of unknown cause, and changes in mood, behaviour or thinking, suicidal thoughts. Inform named consultant urgently.	GP to monitor blood pressure, heart rate. Inform the named consultant of any reported adverse effects i.e. rash, headache urgently.	Specialist: Subject to response to treatment: 6 monthly or 12 monthly if well controlled and stable. Send a letter/results notification to the GP after each clinic attendance indicating current dose, most recent blood pressure, Epworth Sleep Score and frequency of visits. Advise GP on review, duration and or discontinuation of treatment when necessary. Inform GP of patients who do not attend clinic appointments. Advise GP to continue current plan until next reviewed in clinic. GP Request patient seen earlier if condition deterioration or adverse effects experienced between appointments.

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Modafinil

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

- 1. Adverse Effects Patients should be advised to stop treatment if any sign of a rash occurs, any changes in mood, behaviour or thinking or you develop a fast heartbeat, chest pain or unexplained breathlessness. The side-effects mentioned above usually occur in the first 8 weeks of treatment and will be discussed by the doctor initiating the medication.
- Pregnancy and Breast Feeding Modafinil is not recommended for use in pregnancy or breastfeeding. Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.
- 3. Effects on ability to drive or use machinery

Modafinil can cause blurred vision or dizziness in some people therefore it may affect their ability to drive or use machines. Sleepiness associated with condition may also add to this.

Abuse, misuse and diversion

Patients should be carefully monitored for the risk of diversion, misuse and abuse of modafinil.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list) - https://www.medicines.org.uk/emc/product/4319/smpc

Modafinil is a weak enzyme inducer.

Hormonal contraceptives, including oral contraceptive pills, implants, intrauterine contraceptive devices (IUCDs) and contraceptive patches may be less effective when used with modafinil and therefore are not recommended. Advise to use additional barrier methods.

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-agreements/

Methylphenidate (off-label use) - refer to pathway for place in therapy (see links on page 5)

Methylphenidate belongs to a group of medicines called stimulants. Methylphenidate works by stimulating the brain to increase alertness and reduce excessive sleepiness during the day. Although there is no cure for narcolepsy, methylphenidate can help to control symptoms.

Route, Dose, Duration	Monitoring Undertaken by Specialist before requesting shared care	Ongoing monitoring to be undertaken by GP	Stopping Criteria	Monitoring following dose changes	Follow Up
Oral:	Baseline	Check weight, blood	Failure to respond to treatment	GP to monitor blood pressure,	Specialist:
Methylphenidate XL	Blood pressure, heart rate,	pressure and pulse	or adverse effects necessitating	heart rate, weight.	Subject to response to treatment: 6
18-72mg every morning	ECG, Epworth Sleep Score	annually.	withdrawal.	, ,	monthly or 12 monthly if well
Methylphenidate IR	Weight – to allow assessment	,	/	Inform the named consultant	controlled and stable.
10-60mg/day in divided doses.	of weight loss, pregnancy test	Adverse reactions and	Pregnancy	of any reported adverse effects	
	if necessary	inform the named		such as chest pain, fast	Send a letter/results notification to
Renal or hepatic insufficiency	·	consultant if concerns that	Patient request	heartbeat and mood changes	the GP after each clinic attendance
There is no experience with the use of	Ongoing at 6 and 12 monthly	the patient may be	/	urgently.	indicating current dose and Epworth
methylphenidate in patients with renal or	clinic appointments	misusing the medication.	STOP methylphenidate if		sleep score and frequency of visits.
hepatic insufficiency.	Blood pressure, pulse, weight,		changes in mood, behaviour or		Advise GP on review, duration and or
	Epworth Sleep Score		thinking or onset of suicidal		discontinuation of treatment when
Older people	Ask patient about any changes		thoughts.		necessary. Inform GP of patients who
Methylphenidate should not be used in the	to mood, behaviour or		Inform named consultant		do not attend clinic appointments.
elderly. Safety and efficacy has not been	thinking.	/	urgently.		Advise GP to continue current plan
established in this age group.	Ask patient about any chest				until next reviewed in clinic.
	pain, fast heartbeat or				
Duration of Treatment	unexplained breathlessness				GP:
Indefinitely if patient is responding well to	Ask patient about any suicidal				Request patient seen earlier if
treatment and in absence of significant side	thoughts or thoughts about				condition deteriorates or adverse
effects	harming yourself.				effects experienced between
If paradoxical aggravation of symptoms or					appointments.
other serious adverse events occur, the					
dosage should be reduced or discontinued.					

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Adverse effects

- Changes to mood, behaviour or thinking
- Fast heartbeat, palpitations
- Unexplained breathlessness
- Weight loss

2. Pregnancy and Lactation

• Not recommended during pregnancy.

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: SEL Integrated Care 14 Board (covering the boroughs of Bexley/Bromley/Greenwich/ Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-agreements/

• Breast-feeding is not recommended when using Methylphenidate

Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

3. Contra-indications:

The list of contraindications is not exhaustive, please refer to the SPC for detail - https://www.medicines.org.uk/emc/product/6872/smpc

4. Choice of formulation

The choice of formulation of methylphenidate-containing product will be decided by the treating specialist on an individual basis and depends on the intended duration of effect. Generic prescribing permitted.

SPS guidance on prescribing and switching between modified-release methylphenidate preparations can be accessed via this link - https://www.sps.nhs.uk/articles/prescribing-and-switching-between-modified-release-methylphenidate/

5. Abuse, misuse and diversion

• Patients should be carefully monitored for the risk of diversion, misuse and abuse of methylphenidate.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list) - https://www.medicines.org.uk/emc/product/6872/smpc

• Coumarins: Methylphenidate can possibly enhance the effect of coumarins

Dexamfetamine - refer to pathway for place in therapy (see links on page 5)

Dexamfetamine belongs to a group of medicines called stimulants. Dexamfetamine works by stimulating the brain to increase alertness and reduce excessive sleepiness during the day. Although there is no cure for narcolepsy, dexamfetamine can help to control symptoms.

Route, Dose, Duration	Monitoring Undertaken by	Ongoing monitoring to	Stopping Criteria	Monitoring following dose changes	Follow Up
	Specialist before requesting	be undertaken by GP			
	shared care				
Oral: Dexamfetamine 10-60mg daily in divided doses. Renal or hepatic insufficiency There is no experience with the use of dexamfetamine in these patients. Peak plasma levels could be higher and elimination could be prolonged. Dexamfetamine should be used with special caution in this patient group by taking care of titration and dosage.	Baseline Blood pressure, heart rate, ECG, Epworth Sleep Score Weight – to allow assessment of weight loss, pregnancy test if necessary Ongoing at 6 and 12 monthly clinic appointments Blood pressure, pulse, weight, Epworth Sleep Score Ask patient about any changes to mood, behaviour or thinking.	Check weight, blood pressure and heart rate annually. Adverse reactions and inform the named consultant if concerns that the patient may be misusing the medication.	Failure to respond to treatment or adverse effects necessitating withdrawal. Pregnancy Patient request STOP dexamfetamine if changes in mood,	GP to monitor blood pressure, heart rate, weight. Inform the named consultant of any reported adverse effects such as chest pain, fast heartbeat and mood changes urgently.	Specialist: Subject to response to treatment: 6 monthly or 12 monthly if well controlled and stable. Send a letter/results notification to the GP after each clinic attendance indicating current dose and Epworth sleep score and frequency of visits. Advise GP on review, duration and or discontinuation of treatment when necessary. Inform GP of patients who do not attend clinic
Older people	to mood, behaviour or trilliking.		behaviour or thinking or onset		appointments. Advise GP to
			tilliking of offset		

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: SEL Integrated Care 15 Board (covering the boroughs of Bexley/Bromley/Greenwich/Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Dexamfetamine should not be used in the elderly.	Ask patient about any chest	of suicidal	continue current plan until next
Safety and efficacy of dexamfetamine has not been	pain, fast heartbeat or	thoughts. Inform	reviewed in clinic.
established in this age group.	unexplained breathlessness	named consultant	
	Ask patient about any suicidal	urgently.	GP:
Duration of Treatment	thoughts or thoughts about		Request patient seen earlier if
Indefinitely if patient is responding well to treatment	harming yourself.		condition deteriorates or adverse
and in absence of significant side effects.			effects experienced between
If paradoxical aggravation of symptoms or other			appointments.
serious adverse events occur, the dosage should be			
reduced or discontinued.			

Dexamfetamine (cont)

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Adverse effects

- Changes to mood, behaviour or thinking
- Fast heartbeat, palpitations
- Unexplained breathlessness
- Weight loss

2. Pregnancy and Lactation

- Not recommended during pregnancy.
- Breast-feeding is not recommended when using dexamfetamine

Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

3. Contra-indications:

The list of contraindications is not exhaustive, please refer to the SPC for detail.

http://www.medicines.org.uk/emc/medicine/31119

4. Abuse, misuse and diversion

• Patients should be carefully monitored for the risk of diversion, misuse and abuse of dexamfetamine.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list) - http://www.medicines.org.uk/emc/medicine/31119

MAO-A and MAO-B inhibitors: Risk of hypertensive crisis if given with dexamfetamine

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Venlafaxine XL/IR (off-label use)- refer to pathway for place in therapy (see links on page 5)

Venlafaxine belongs to a group of medicines called anti-depressants. It can be used to treat cataplexy in narcolepsy. Venlafaxine is thought to work by interfering with certain

Route, Dose, Duration	Monitoring Undertaken by Specialist before requesting shared care	Ongoing monitoring to be undertaken by GP	Stopping Criteria	Monitoring following dose changes	Follow Up
Oral: Venlafaxine XL 37.5-150mg in the morning.	Baseline Blood pressure, heart	Blood pressure and heart rate	Failure to respond to treatment or adverse effects	GP to monitor blood pressure and heart	Specialist: Subject to response to treatment: 6 monthly or 12
Venlafaxine IR 37.5-150mg daily (to confirm with sleep specialist if once or twice daily dosing	rate, cataplexy severity, Epworth Sleep	annually.	necessitating withdrawal.	rate.	monthly if well controlled and stable.
required)	Score.		Patient request	Inform the named consultant of any	Send a letter/results notification to the GP after each clinic attendance indicating current dose and Epworth
Duration of Treatment Indefinitely if patient is responding well to treatment	Ongoing at 6 and 12 monthly clinic		Inform named consultant urgently. Sleep centre will	reported adverse effects such as	sleep score and frequency of visits.
and in absence of significant side effects.	appointments		advise appropriate weaning regimen.	hypertension, anxiety urgently.	Advise GP on review, duration and or discontinuation of treatment when necessary. Inform GP of patients
Abrupt cessation: Abrupt cessation of antidepressants can result in	Blood pressure, heart rate, cataplectic			,	who do not attend clinic appointments. Advise GP to continue current plan until next reviewed in clinic.
status-cataplecticus (particularly cessation of venlafaxine). They should be withdrawn slowly and	episodes, Epworth Sleep Score.				·
overlapped with next treatment option.					GP: Request patient seen earlier if condition deterioration
Hepatic impairment 50% dose reduction should be considered mild and					or adverse effects experienced between appointments
moderate hepatic impairment. Risk versus benefit in patients with severe hepatic impairment.					
Renal impairment No change in dosage if caution is advised. Dose	/				
should be reduced by 50% in haemodialysis and in severe renal impairment (GFR < 30 ml/min).					
Older people: No dose adjustment required. Caution should be					
exercised using lowest effective dose. Careful					
monitoring is required when an increase in the dose is required.					

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: SEL Integrated Care 17 Board (covering the boroughs of Bexley/Bromley/Greenwich/ Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Venlafaxine (cont)

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Choice of formulation

Modified release preparation should ONLY be prescribed for the management of cataplexy. Using the immediate release preparation may increase the risk of rebound cataplexy or the onset of status cataplecticus.

2. Adverse effects

- Abnormal dreams, insomnia,
- Anxiety; particularly on withdrawal
- Dizziness, drowsiness, confusion, nervousness
- Sweating, nausea & vomiting
- Hypertension, palpitations

3. Effects on Ability to Drive and use Machines

• Any psychoactive medicinal product may impair judgment, thinking, and motor skills. Therefore, any patient receiving venlafaxine should be cautioned about their ability to drive or operate hazardous machinery.

4. Contraindications

• Conditions associated with high risk of cardiac arrhythmia; uncontrolled hypertension

5. Pregnancy and Lactation

- Avoid unless potential benefit outweighs risk. Risk of withdrawal effects in neonate.
- Present in breast milk- avoid
- Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list)

The list of significant drug interactions is not exhaustive, please refer to the SPC for detail.

https://www.medicines.org.uk/emc/product/2686/smpc

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Clomipramine - refer to pathway for place in therapy (see links on page 5)

Clomipramine belongs to a group of medicines called anti-depressants. It can be used to treat cataplexy in narcolepsy. Clomipramine works by interfering with certain chemicals in

the brain which may be involved in causing the symptoms of cataplexy.

the brain which may be involved in causing the symptoms of cataplexy.								
Route, Dose, Duration	Monitoring Undertaken by	Ongoing monitoring	Stopping Criteria	Monitoring following	Follow Up			
	Specialist before requesting	to be undertaken by		dose changes				
	shared care	GP						
Oral:	Baseline	Blood pressure and	Failure to respond to	GP to monitor blood	Specialist:			
Clomipramine 10-75mg at night.	Blood pressure, heart rate,	heart rate annually.	treatment or adverse effects	pressure and heart rate.	Subject to response to treatment: 3			
Duration of Treatment	cataplexy severity, Epworth Sleep		such as hallucinations,		monthly, 6 monthly or 12 monthly if			
Indefinitely if patient is responding well to treatment	Score.		disorientation, agitation or	Inform the named	well controlled and stable.			
and in absence of significant side effects.	Ongoing at 6 and 12 monthly		anxiety necessitating	consultant of any	6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			
	clinic appointments		withdrawal.	reported adverse effects such as	Send a letter/results notification to the GP after each clinic attendance			
Treatment cessation			Patient request	hallucinations,	indicating current dose and Epworth			
Withdrawal effects such as nausea, vomiting,	Blood pressure, heart rate,			disorientation, agitation	sleep score and frequency of visits.			
abdominal pain, diarrhoea, insomnia, headache,	cataplectic episodes, Epworth		Inform named consultant	or anxiety urgently.	steep score and mediative, or visits.			
nervousness and anxiety may occur within 5 days of	Sleep Score.	/	urgently. Sleep centre will		Advise GP on review, duration and or			
stopping treatment. May be mild and self-limiting, can be severe. The risk of withdrawal symptoms is			advise appropriate weaning		discontinuation of treatment when			
increased if treatment stopped suddenly after regular			regimen.		necessary. Inform GP of patients			
administration for \geq 6 weeks. Reduce gradually over 4					who do not attend clinic			
weeks, or longer if withdrawal symptoms emerge					appointments. Advise GP to continue			
Tricyclic and related antidepressants should be					current plan until next reviewed in			
withdrawn slowly.					clinic.			
Renal impairment	/							
Use with caution					GP:			
					Request patient seen earlier if condition deterioration or adverse			
Hepatic impairment					effects experienced between			
Sedative effects are increased in hepatic impairment.	/				appointments.			
Avoid in severe liver disease.					арропилено.			
Older people								
May show a stronger response to clomipramine. Use								
with caution. Doses should be increased cautiously.								
The sacretic boses should be moreused sautiously.								

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Clomipramine (cont)

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Adverse effects

- Abdominal pain, constipation (diarrhoea associated with withdrawal)
- Fatigue, impaired memory
- Aggression, restlessness
- Hypertension
- Flushing, muscle weakness/twitching

2. Effects on Ability to Drive and use Machines

• Drowsiness may affect the performance of skilled tasks (e.g. driving).

3. Contraindications

- Arrhythmias
- Acute porphyrias
- during the manic phase of bipolar disorder
- major depression
- heart block
- immediate recovery period after myocardial infarction

4. Pregnancy and Lactation

- Clomipramine is not recommended for use in women not using contraception
- Neonatal withdrawal symptoms reported if used during third trimester
- The quantity of clomipramine secreted into breast milk is small. Nursing mothers should be advised to withdraw from the medication or cease breast-feeding.
- Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

5. Anticholinergic burden

Elderly patients are particularly sensitive to anticholinergic, neurological, psychiatric, or cardiovascular effects. Their ability to metabolise and eliminate drugs may be reduced, leading to a risk of elevated plasma concentrations at therapeutic doses. Dose range in pathway is appropriate for older people (10-75mg daily)

Clinically Significant Drug Interactions (refer to BNF/SPC for full list)

The list of significant drug interactions is not exhaustive, please refer to the SPC for detail.

https://www.medicines.org.uk/emc/product/2550

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Fluoxetine (off-label use)- refer to pathway for place in therapy (see links on page 5)

Fluoxetine belongs to a group of medicines called anti-depressants. It can be used to treat cataplexy in narcolepsy. Fluoxetine works by interfering with certain chemicals in the

Route, Dose, Duration	Monitoring Undertaken by Specialist before	Ongoing monitoring to be undertaken by GP	Stopping Criteria	Monitoring following dose changes	Follow Up
Oral:	Baseline	Blood pressure and heart rate	Failure to respond to treatment or adverse	GP to monitor blood pressure and heart rate	Specialist: Subject to response to treatment: 6 monthly or
Fluoxetine 20-60mg daily in the morning.	Blood pressure, heart rate, cataplexy severity, Epworth Sleep Score.	annually.	effects necessitating withdrawal.	for patients on concomitant medications that may prolong QT or for those already under hypertension management.	12 monthly if well controlled and stable.
Hepatic impairment Reduce dose or increase dosing interval.	Weight – to allow assessment of weight		Patient request	Inform the named consultant of any reported adverse effects such as	Send a letter/results notification to the GP after each clinic attendance indicating current dose and Epworth sleep score and frequency of visits.
Older people	loss		Inform named consultant	palpitations, anxiety, urinary	
Caution is recommended when increasing the dose. Daily dose should generally not exceed 40 mg. Maximum recommended dose is 60 mg/day.	Ongoing at 6 and 12 monthly clinic appointments		urgently. Sleep centre will advise appropriate weaning regimen.	frequency/retention, weight loss urgently.	Advise GP on review, duration and or discontinuation of treatment when necessary. Inform GP of patients who do not attend clinic appointments. Advise GP to continue current plan until next reviewed in clinic.
Duration of Treatment Indefinitely if patient is responding well to treatment and in absence of significant side effects. If paradoxical aggravation of symptoms or other serious adverse events occur, the dosage should be reduced or discontinued.	Blood pressure, heart rate, cataplectic episodes, Epworth Sleep Score.				GP: Request patient seen earlier if condition deterioration or adverse effects experienced between appointments.

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

Fluoxetine (cont)

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Adverse effects

- Abdominal pain, diarrhoea, constipation
- Nausea & vomiting, GI effects, dyspepsia
- Dizziness, drowsiness, visual disturbances, hallucinations
- Urinary retention, sexual dysfunction
- · Bleeding disorders
- Anxiety, nervousness

2. Effects on Ability to Drive and use Machines

• May also impair performance of skilled tasks (e.g. driving, operating machinery)

3. Contraindications

The list of contraindications is not exhaustive, please refer to the SPC for detail- https://www.medicines.org.uk/emc/product/15589/smpc

4. Pregnancy and Lactation

Manufacturers advise to avoid during pregnancy unless the potential benefit outweighs the risk. There is a small increased risk of congenital heart defects when taken during early pregnancy. If used during the third trimester there is a risk of neonatal withdrawal symptoms, and persistent pulmonary hypertension in the newborn has been reported.

Present in breast milk- avoid

Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list)

The list of significant drug interactions is not exhaustive, please refer to the SPC for detail - https://www.medicines.org.uk/emc/product/15589/smpc

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-im

Sodium Oxybate - refer to pathway for place in therapy (see links on page 5)

Sodium Oxybate is used to treat narcolepsy when cataplexy is also a problem. Sodium Oxybate promotes deep sleep and improves night-time sleep. It helps with excessive daytime sleepiness as well as helping the symptoms of cataplexy. It is sometimes used in combination with other medicines for narcolepsy. Although there is no cure for narcolepsy, sodium oxybate can help to control symptoms.

Route, Dose, Duration	Monitoring Undertaken by Specialist	Ongoing monitoring to	Stopping Criteria	Monitoring following	Follow Up
	before requesting shared care	be undertaken by GP		dose changes	
Oral:	Baseline	Check weight, blood	Failure to respond to	Check weight, blood	Specialist:
The recommended starting dose is 4.5 g/day sodium	Blood pressure, heart rate, Epworth	pressure and pulse	treatment or adverse	pressure and pulse	Subject to response to treatment: 6
oxybate (9 ml Xyrem®) divided into two equal doses of	Sleep Score, pregnancy test if	annually.	effects necessitating	annually.	monthly or 12 monthly if well
2.25 g/dose (4.5 ml/dose).	necessary, weight – to allow		withdrawal.		controlled and stable.
The dose should be titrated to effect based on efficacy	assessment of weight loss, severity of	Adverse reactions and			
and tolerability up to a maximum of 9 g/day divided into	cataplectic episodes. oximetry (only	inform the named	Pregnancy	Inform the named	Send a letter/results notification to
two equal doses	avoid in severe OSA) mood	consultant if concerns		consultant of any	the GP after each clinic attendance
of 4.5g/dose (9ml/dose) by adjusting up or down in dose		that the patient may be	Patient request	reported adverse effects	indicating current dose and
increments of 1.5 g/day (i.e. 0.75 g/dose or 1.5 ml/dose).		misusing the		such as sleep walking,	Epworth sleep score and frequency
A minimum of two weeks is recommended between	Ongoing at 6 and 12 monthly clinic	medication.	STOP sodium oxybate if	nocturnal enuresis	of visits.
dosage increments.	appointments		changes in mood,	urgently.	
 Each dose of Xyrem® must be diluted with 60 ml of 		/	behaviour or thinking		Advise GP on review, duration and
water in the dosing cup prior to ingestion. Single doses	Blood pressure, heart rate, Epworth		or onset of suicidal		or discontinuation of treatment
of 4.5g should not be given unless the patient has been	Sleep Score, pregnancy test if		thoughts. Inform		when necessary. Inform GP of
titrated previously to that dose level.	necessary, weight – to allow		named consultant		patients who do not attend clinic
Because food significantly reduces the bioavailability of	assessment of weight loss.		urgently.		appointments. Advice GP to
sodium oxybate, patients should eat at least several (2-	Cataplexy epsiodes.				continue current plan until next
3) hours before taking the first dose of Xyrem® at	Ask the patient about any changes in		Sleep centre will advise		reviewed in clinic.
bedtime. Patients should always observe the same	mood, behaviour or side effects such		an appropriate weaning		
timing of dosing in relation to meals. Xyrem® should be	as nocturnal enuresis, sleep walking.		regimen.		GP:
taken orally upon getting into bed and again between	Any change in breathing i.e increased				Request patient seen earlier if
2.5 to 4 hours later.	snoring, reports of apnoea.				condition deterioration or adverse
Patients advised to set an alarm and to remain in bed	/				effects experienced between
while they take their second dose.					appointments.
Discontinuation of Xyrem®: The discontinuation effects	/				
of sodium					
oxybate have not been systematically evaluated in					
controlled clinical trials. If the patient stops medication					
for more than 14 consecutive days, titration should be					
restarted as per initiation regimen.					

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: SEL Integrated Care 23 Board (covering the boroughs of Bexley/Bromley/Greenwich/ Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/sel-imoc-shared-care-agreements/

Patients with hepatic impairment: The starting dose should be halved in patients with hepatic impairment, and response to dose increments			
monitored closely. Patients with renal impairment: Patients with impaired			
renal function should consider a dietary recommendation to reduce sodium intake.			
Elderly patients: Elderly patients should be monitored closely for impaired motor and/or cognitive function when taking sodium oxybate.			
Duration of Treatment: Indefinitely if patient is responding well to treatment and in absence of significant side effects.			

Sodium Oxybate (cont)

Practical issues including adverse effects, interactions, other relevant advice and information (refer to BNF/SPC for full list):

1. Storage

Sodium oxybate comes in a bottle with two containers for mixing the medication. Both the bottle and storage containers have child proofed caps. The mixed solution should be kept in a bed side drawer just before going to sleep. If there are children in the house, this draw should be locked or be out of the reach of children. Discard diluted solution after 24 hours.

2. Adverse effects

- Sleep walking
- Urinary incontinence
- Drowsiness, dizziness, blurred vision
- Nausea, vomiting, weight loss, abdominal pain
- Hypertension, palpitations

3. Effects on Ability to Drive and use Machines

- Sodium oxybate has a major effect on the ability to drive and use machines. These patients should already be known to the DVLA
- For at least 6 hours after taking sodium oxybate, patients must not undertake activities requiring complete mental alertness or motor coordination, such as operating machinery or driving.
- When patients first start taking sodium oxybate, they should take extreme care when driving, operating heavy machines or performing any other task which is dangerous or requires full mental alertness.

4. Contraindications

The list of contraindications is not exhaustive, please refer to the SPC for detail.

https://www.medicines.org.uk/emc/product/178

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: SEL Integra

South East London Shared Care Prescribing Guideline for the management Narcolepsy (+/- Cataplexy) and

Idiopathic Hypersomnia in adults – stimulant therapy and anti-cataplectic agents

Original Approval Date: August 2018 Last Reviewed and updated: October 2025 Review date: October 2028 (or sooner if evidence or practice changes)

Document control: If this transfer of prescribing responsibility form is printed or downloaded it becomes uncontrolled. The latest approved version can be accessed via: <a href="https://www.selondonics.org/icb/healthcare-professionals/medicines/sel-imoc/s

5. Pregnancy and Lactation

- Not recommended during pregnancy.
- Breast-feeding is not recommended when using Sodium oxybate.

Any patient considering becoming pregnant or has discovered they are pregnant should be referred back to their consultant immediately and shared care will no longer apply for the duration of the pregnancy and while breast feeding.

Clinically Significant Drug Interactions (refer to BNF/SPC for full list) - https://www.medicines.org.uk/emc/product/178

- Alcohol: The combined use of alcohol with sodium oxybate may result in potentiation of the central nervous system-depressant effects of sodium oxybate.
- Sedative hypnotics or other CNS depressants: Sodium oxybate should not be used in combination
- Since sodium oxybate is metabolised by GHB dehydrogenase there is a potential risk of an interaction with drugs that stimulate or inhibit this enzyme (e.g. valproate, phenytoin or ethosuximide). No interaction studies have been conducted in human subjects, although if sodium oxybate and valproate are used concomitantly, a decrease in sodium oxybate dose by 20% is recommended.
- Possible additive effect of antidepressants and sodium oxybate cannot be excluded. The rates of adverse events are increased when sodium oxybate is co-administered with tricyclic antidepressants.
- There is a higher risk of sleep apnoea in patients with BMI ≥ 40kg/m2.